



Distal Radial Fractures Treated with Volar ORIF AGGRESSIVE PROTOCOL

Jennifer L. Cook, M.D.
Board Certified
Orthopaedic Surgery

Stephen A. Hanff, M.D.
Board Certified
Orthopaedic Surgery

Aaron K. Mates, M.D.
Board Certified
Orthopaedic Surgery

Troy D. Pashuck, M.D.
Orthopaedic Surgery

- **DAYS THREE – FIVE:**

- Finger, Elbow, & Shoulder Active Range Of Motion
- Wrist & Forearm Active Range Of Motion
- Finger Passive Range of Motion
- Edema Control
- Custom Splint (30° Wrist Extension)
 - Splint may be Removed for Hygiene, dressing, eating and exercises

- **TWO WEEKS:**

- Wrist & Forearm Passive Range of Motion
- Isometrics
- Light Putty Strengthening

- **THREE WEEKS:**

- Weaned from Splint

- **FOUR WEEKS:**

- Isotonic Strengthening
- Medium Putty Strengthening
- Discontinuation of Splint Use

- **SIX WEEKS:**

- Heavy Putty Stengthening

- **TWELVE WEEKS:**

- Study Visit For Measurements

- **SIX MONTHS:**

- Study Visit For Measurements

A REGISTERED LIMITED LIABILITY PARTNERSHIP FOR ORTHOPAEDIC SURGERY

2165 Little Road • Trinity, FL 34655 • 727-37-BONES (727-372-6637) • Fax 727-375-5044 • www.floridajointcare.com