



Rotator Cuff Type III Repair (Large Tear)

This rehabilitation protocol has been developed for a patient following a Rotator Cuff Type III Repair. It is extremely important to **protect the repair for 6 weeks** post – operatively to allow for appropriate healing to take place. This protocol has been divided into several phases, each of which may vary slightly based upon the individual patient and his or her special circumstances.

Jennifer L. Cook, M.D.
Board Certified
Orthopaedic Surgery

Stephen A. Hanff, M.D.
Board Certified
Orthopaedic Surgery

Aaron K. Mates, M.D.
Board Certified
Orthopaedic Surgery

Troy D. Pashuck, M.D.
Orthopaedic Surgery

Goals of the Surgical Procedure and Rehabilitation:

- Control Pain and Inflammation
- Maintain the Integrity of the repair
- Prevent Muscular Inhibition
- Regain normal upper extremity strength and endurance
- Regain normal shoulder range of motion
- Achieve the highest level of function based on both the physician and the patient's individual goals.

Important Post – Operative Signs to Monitor:

- Swelling of the shoulder and surrounding soft tissue
- Abnormal pain response, hypersensitive – an increase in night pain
- Severe Range of Motion Limitations
- Weakness in the upper extremity musculature

Returning to Activity:

*To safely and most efficiently return to normal or high level functional activity the patient requires adequate strength, flexibility, and endurance this **will require both time and clinical evaluation.** Functional evaluation including strength and range of motion testing is one method of evaluating a patient's readiness to return to such activity. Returning to intense activities following a Rotator Cuff Type III Repair will require both a strenuous strengthening and range of motion program. The patient will also have to wait a period of time to allow for appropriate tissue healing. Symptoms such as pain, swelling, or instability should be closely monitored by the patient.*

PHASE I: IMMEDIATE POST – SURGICAL PHASE (DAY 1 – WEEK 8)

Goals:

- Maintain integrity of the Repair
- Gradually increase Passive Range of Motion
- Diminish Pain and Inflammation
- Prevent Muscular Inhibition

A REGISTERED LIMITED LIABILITY PARTNERSHIP FOR ORTHOPAEDIC SURGERY

2165 Little Road • Trinity, FL 34655 • 727-37-BONES (727-372-6637) • Fax 727-375-5044 • www.floridajointcare.com

Day One to Six Weeks:

- Use Abduction Pillow Brace
- Cryotherapy for Pain and Inflammation
 - Ice 15 – 20 minutes every hour
- Sleeping
 - Sleep in Pillow Brace

Week 6 to Week 8:

- Discontinue use of the pillow brace
- Pendulum Exercises
- Passive ROM to Tolerance
 - Flexion to at least 115 degrees
 - ER in Scapular Plane at 45 degrees of Abduction to 20 – 25 degrees
 - IR in scapular plane at 45 degrees abduction to 30 – 35 degrees
- Active Assisted ROM Exercises (Wand Exercises)
 - ER / IR in scapular plane at 45 degrees abduction
 - Flexion to Tolerance
 - Therapist should provide assistance by supporting the patient's arm (Especially when arm is lowering)
- Elbow / Hand ROM & Gripping Exercises
- Isometrics (Submaximal and Subpainful)
 - Flexion with Elbow Bent
 - Extension with Bent Elbow
 - Abduction with Bent Elbow
 - ER / IR with arm in Scapular Plane
- Gentle Elbow Strengthening Exercises
- Initiate Rhythmic Stabilization ER / IR at 45 degrees abduction
- Use Ice for Pain Control

Precautions:

- No Lifting objects
- No Excessive Shoulder Extension
- No Excessive stretching or Sudden Movements
- No Supporting of body weight by hands

PHASE II: PROTECTION PHASE (WEEKS 8 - 10)

Goals:

- Allow Healing of soft tissue
- Do not overstress healing tissue
- Gradually restore full Passive ROM
- Re-Establish Dynamic Shoulder Stability
- Continue to decrease pain and inflammation

Week 8 to Week 9

- Passive Range of Motion to tolerance
 - Flexion to 140 – 155 degrees
 - ER at 90 degrees abduction to at least 45 degrees
 - IR at 90 degrees abduction to at least 45 degrees
- Active Assisted ROM to tolerance
 - Flexion (continue to support arm)
 - ER/IR in scapular plane at 45 degrees abduction
 - ER / IR at 90 degrees Abduction
- Dynamic Stabilization Drills
 - Rhythmic Stabilization Drills
 - ER/IR in Scapular Plane
 - Flexion / Extension at 100 degrees Flexion and 125 degrees Flexion
- Continue all Isometric Contractions
- Initiate Scapular Isometrics
- Initiate ER/IR strengthening using exercise tubing at 0 degrees of abduction using light resistance and use towel roll under arm
- Initiate light manual resistance ER supine in Scapular plane
- Initiate Prone Rowing and Prone Shoulder Extension
- Initiate ER strengthening exercises
- Initiate Isotonic Elbow Flexion
- Continue all Precautions
 - No Lifting
 - No Excessive Motion
- May continue to use Cryotherapy as needed
- May use pool for light AROM exercises
- Patient should exhibit full PROM by week 8

Week 9 to Week 10

- May use heat prior to exercises
- Continue AAROM and Stretching exercises
 - Especially for movements that are not full ROM
 - Shoulder Flexion
 - ER at 90 degrees Abduction
- Initiate AROM exercises
 - Shoulder flexion scapular plane
 - Shoulder abduction
- Progress Isotonic Strengthening Exercise Program
 - ER Tubing
 - Side-lying IR
 - Prone Rowing
 - Prone Horizontal Abduction
 - Bicep Curls

Precautions:

- No heavy lifting of objects
- No excessive behind the back reaching
- No supporting of body weight by hands and arms
- No sudden jerking Motions

PHASE III: INTERMEDIATE PHASE (WEEKS 10 – 14)

GOALS:

- Full AROM
- Maintain Full Passive ROM
- Dynamic Shoulder Stability
- Gradual Restoration of shoulder strength
- Gradual Return To Functional Activities

Week 10 to Week 14

- Continue stretching and PROM as needed to maintain Full ROM
- Continue Dynamic Stabilization Drills
- Progress Strengthening Program
 - ER/IR Tubing
 - ER Side-lying
 - Lateral Raises
 - Full Can In Scapular Plane
 - Prone: Rowing / Horizontal Abduction / Shoulder Extension
 - Elbow Flexion / Extension
 - If Physician Permits – May Initiate LIGHT Functional Activities
 - Progress to fundamental Shoulder Exercises
 - Therapist may initiate isotonic resistance 1# during flexion and abduction
- ***if non painful normal motion is exhibited***

****Progress patient strengthening program (Increase 1 lb. / 10 days) if and only if...Patient is able to elevate arm without shoulder or scapular hiking, if un-capable, continue with glenohumeral joint exercises without increasing difficulty****

PHASE IV: ADVANCED STRENGTHENING PHASE (Weeks 14 – 22)

Goals:

- Maintain Full Non Painful ROM
- Enhance Functional Use of UE
- Improve Muscular Strength and Power
- Gradual Return to Functional Activities

Week 15

- Continue ROM & Stretching to Maintain Full ROM

A REGISTERED LIMITED LIABILITY PARTNERSHIP FOR ORTHOPAEDIC SURGERY

2165 Little Road • Trinity, FL 34655 • 727-37-BONES (727-372-6637) • Fax 727-375-5044 • www.floridajointcare.com

- Self-Capsular Stretches
- Progress Shoulder Strengthening Exercises
 - Fundamental Shoulder Exercises
- Initiate Interval Golf Program (If Appropriate)

Weeks 20 – 22

- Continue all Exercises above
- Progress Golf Program to Playing Golf (if Appropriate)
- Initiate Interval Tennis Program (If Appropriate)
- May Initiate Swimming

PHASE IV: RETURN TO ACTIVITY PHASE (WEEKS 23 – 26)

Goals:

- Gradual Return to Strenuous Work Activities
- Gradual Return to Recreational Sport Activities

Week 23 ---- on

- Continue Fundamental Shoulder Exercise Program (At Least 4 Times Per Week)
- Continue Stretching, If Motion is Tight
- Continue Progression to Sport Participation