



TOTAL KNEE ARTHROPLASTY PROTOCOL

PHASE 1: IMMEDIATE POST SURGICAL PHASE (DAY 0 – 10)

Goals:

- Achieve Quad Contraction
- Safe Independent Ambulation with Walker or crutches as needed
- Passive knee extension to 0 degrees
- Knee Flexion to 90 degrees or greater
- Control:
 - Inflammation
 - Swelling
 - Bleeding

Jennifer L. Cook, M.D.
Board Certified
Orthopaedic Surgery

Stephen A. Hanff, M.D.
Board Certified
Orthopaedic Surgery

Aaron K. Mates, M.D.
Board Certified
Orthopaedic Surgery

Troy D. Pashuck, M.D.
Orthopaedic Surgery

Day 0 - 2:

- Weight bearing as tolerated (*Unless otherwise Specified by Physician*)
 - Walker
 - 2 crutches (*If Balance is Sufficient*)
- Cryotherapy
 - *Immediately and continuously*
 - unless ambulating
- ROM of knee to begin immediately post operatively
- Exercises:
 - Ankle Pumps
 - Passive knee extension to 0 degrees
 - Straight Leg Raises
 - Quad Sets
 - Knee Flexion to at least 90 degrees
 - Knee Extension to 0 degrees
 - Instruct:
 - Gait Training
 - Safe Transfers

Day 3 – 10

- Weight Bearing as tolerated
 - Walker
 - Two crutches
- Cryotherapy
- Exercises

A REGISTERED LIMITED LIABILITY PARTNERSHIP FOR ORTHOPAEDIC SURGERY

2165 Little Road • Trinity, FL 34655 • 727-37-BONES (727-372-6637) • Fax 727-375-5044 • www.floridajointcare.com

- Ankle Pumps
- Passive Knee extension to 0 degrees
- Straight Leg Raises
- Quad Sets
- AAROM:
 - Knee Flexion to at least 90 degrees
- Hip Abduction / Adduction
- Instruct
 - Gait Training
 - Safe Transfers
- Start Stationary Bike
 - Low Resistance

PHASE II: MOTION PHASE (WEEK 2 – 6)

Goals

- Improve ROM
- Enhance Muscular Strength and endurance
- Dynamic Joint Stability
- Diminish Swelling / Inflammation
- Establish / Return to functional Activities

Criteria to enter Phase II

- Leg Control
 - Able to perform Straight Leg Raise
- Active ROM
 - 0 - 90 degrees
- Minimal Pain / Swelling
- Independent
 - Ambulation
 - Transfers

Weeks 2 – 4

- Weight Bearing with Assistive device as needed
- Wean from a walker to a cane OR from 2 crutches to 1 crutch **by 2 weeks**
- Wean off Assistive Device **Completely by no later than 4 weeks**
- Exercises
 - Quad Sets
 - VMO Recruitment
 - Straight Leg Raises
 - VMO Recruitment
 - Knee Extension 90 – 0 degrees
 - Terminal Knee Extension 45 – 0 degrees
 - Hip Abduction / Adduction
 - Hamstring curls

- Knee Flexion to at least 115 degrees
- Stretches
 - Hamstrings
 - Gastrocnemius
 - Soleus
 - Quads
 - Passive Knee Extension Stretch
- Continue Stationary bike
 - Advance Resistance as tolerated
- Continue with Cryotherapy
- Patellofemoral Mobilization
- Incision Mobilization
- Patients may begin to drive
 - **IF** no longer using assistive devices for ambulation
 - *Approx. 2 weeks post op*

Weeks 4 – 6

- Exercises
 - Continue previous exercises
 - Initiate step ups
 - Front
 - Lateral
 - Advance resistance on stationary bike
 - Initiate progressive walking program
 - Initiate endurance pool programs
 - Swimming with flutter kicks
 - Return to functional Activities

***** Continue compression, ice, elevation as needed for swelling. Patients *should be walking and driving independently* at this point *****

PHASE III: INTERMEDIATE PHASE (WEEKS 7 – 12)

Goals

- Progression of ROM to greater than 115 degrees
- Enhancement of strength and endurance
- Eccentric / Concentric control of Lower Extremity
- Cardiovascular Fitness
- Functional Activity Performance

Criteria To Enter Phase III

- Range of Motion = 0 – 115 Degrees
- Voluntary Quad Control

- Independent Ambulation
- Minimal Pain

Weeks 7 – 12

- Exercises
 - Continue:
 - Previous Exercises
 - Pool Activities
 - Walking
 - Stationary Bike
 - Aggressive AROM 0 – 115 degrees
 - Strengthen Quadriceps / Hamstrings

PHASE IV: ADVANCED ACTIVITY PHASE (WEEKS 12 & BEYOND)

Goals:

- Allow patients to return to advanced level of function
 - Recreational Sports
- Maintain / Improve strength and endurance of lower extremity
- Return to normal life and routine

Criteria to enter Phase IV:

- Full Non Painful ROM 0 – 115 degrees
- Strength 90% of contralateral lower extremity (if that extremity is normal)
- Minimal Pain and swelling
- Satisfactory clinical Examination

Week 12 and Beyond

- Exercises
 - Quad sets
 - Straight leg raises
 - Step ups
 - Hip abduction / adduction
 - Knee extension
 - Stationary bike
 - Swimming
 - Walking
 - Stretching 0 – 115 degrees

***** Return to Pre-operative Activities and Develop HEP to Maintain Leg Function *****

***** NO SQUATS OR LUNGES AT ANY TIME *****